

On q -Runge domains

Youssef ALAOUI

Summary. In [2], Coltoiu gave an example of a domain $D \subset \mathbb{C}^6$ which is 4-complete such that for every $\mathcal{F} \in Coh(\mathbb{C}^6)$ the restriction map $H^3(\mathbb{C}^6, \mathcal{F}) \rightarrow H^3(D, \mathcal{F})$ has a dense image but D is not 4-Runge in $\mathbb{C}^6$. Here, we prove that for every integers $n \geq 4$ and $1 \leq q \leq n$ there exists a domain $D \subset \mathbb{C}^n$ which is not $(\tilde{q} - 1)$ -Runge in $\mathbb{C}^n$ but such that for any coherent analytic sheaf $\mathcal{F}$ on $\mathbb{C}^n$ the restriction map $H^p(\mathbb{C}^n, \mathcal{F}) \rightarrow H^p(D, \mathcal{F})$ has a dense image for all $p \geq \tilde{q} - 2$ if q does not divide n , where $\tilde{q} = n - [\frac{n}{q}] + 1$ and $[\frac{n}{q}]$ denotes the integral part of $\frac{n}{q}$.

Key words: Stein spaces; q -convex functions; q -complete and q -Runge complex spaces.

2000 MS Classification numbers: 32E10, 32E40.

1 Introduction

An open subset D of a q -complete space X is said to be q -Runge in X if for any compact subset $K \subset D$ there exists a q -convex exhaustion function $\phi : X \rightarrow \mathbb{R}$ such that $K \subset \{x \in X : \phi(x) < 0\} \subset\subset D$. When $D \subset X$ is q -Runge in X it follows from an important result of Andreotti and Grauert [1] that for every $\mathcal{F} \in coh(X)$ the restriction map $H^p(X, \mathcal{F}) \rightarrow H^p(D, \mathcal{F})$ has a dense image for all $p \geq q - 1$.

A counterexample to the converse of this statement is given by Coltoiu in [2] where it is shown that there exists a domain $D \subset \mathbb{C}^6$ which is 4-complete but not 4-Runge in $\mathbb{C}^6$ such that for every $\mathcal{F} \in Coh(\mathbb{C}^6)$ the restriction map $H^3(\mathbb{C}^6, \mathcal{F}) \rightarrow H^3(D, \mathcal{F})$ has a dense image.

The purpose of this paper is to prove the following.

theorem 1 -Let (n, q) be a pair of integers with $1 \leq q \leq n$. Suppose that $n \geq 4$ and q does not divide n . Then there is a domain $D \subset \mathbb{C}^n$ which is not $(\tilde{q} - 1)$ -Runge in $\mathbb{C}^n$ but for any coherent analytic sheaf $\mathcal{F}$ on $\mathbb{C}^n$ the restriction map $H^p(\mathbb{C}^n, \mathcal{F}) \rightarrow H^p(D, \mathcal{F})$ has a dense image for all $p \geq \tilde{q} - 2$.

Examples proving theorem 1

Let (n, q) be a pair of integers with $1 \leq q \leq n$. We put $m = \lfloor \frac{n}{q} \rfloor$ and suppose that $r = n - mq > 0$. We consider the functions $\phi_1, \dots, \phi_{m+1}$ defined on $\mathbb{C}^n$ by

$$\phi_j = \sigma_j + \sum_{i=1}^m \sigma_i^2 - \frac{1}{4} \|z\|^2 + N \|z\|^4, \quad j = 1, \dots, m,$$

and

$$\phi_{m+1} = -\sigma_1 - \dots - \sigma_m + \sum_{i=1}^m \sigma_i^2 - \frac{1}{4} \|z\|^2 + N \|z\|^4,$$

where $\sigma_j = y_j + \sum_{i=m+1}^n |z_j|^2 - (m+1) \sum_{i=m+(j-1)(q-1)+1}^{m+j(q-1)} |z_j|^2$ for $j = 1, \dots, m$,

$z_j = x_j + iy_j$. Then it is known from [4] that all ϕ_j , $1 \leq j \leq m+1$, are q -convex on $\mathbb{C}^n$, if $N > 0$ is sufficiently large and, if $\rho = \max\{\phi_j, 1 \leq j \leq m+1\}$, then for $\varepsilon_o > 0$, the set $D_{\varepsilon_o} = \{z \in \mathbb{C}^n : \rho(z) < -\varepsilon_o\}$ is relatively compact in the unit ball $B = B(0, 1)$, if N is large enough and $\varepsilon_o > 0$ is sufficiently small.

lemma 1 -For every coherent analytic sheaf $\mathcal{F}$ on $\mathbb{C}^n$, the cohomology groups $H^p(D_{\varepsilon_o}, \mathcal{F})$ vanish for all $p \geq \tilde{q} - 2$.

Proof

The proof is by induction on m .

We have $D_{\varepsilon_o} = D_1 \cap \dots \cap D_{m+1}$, where $D_i = \{z \in \mathbb{C}^n : \phi_i(z) < -\varepsilon_o\}$ is obviously q -complete and q -Runge in $\mathbb{C}^n$. In fact, $\frac{-1}{\phi_i + \varepsilon_o}$ is a q -convex exhaustion function on D_i and, if $K \subset D_i$ is a compact subset and $\psi \in C^\infty(\mathbb{C}^n, (0, \infty))$ a strictly plurisubharmonic exhaustion function on $\mathbb{C}^n$, then for every real number $C > 0$, the function $\psi_C = \psi + C(\phi_i + \varepsilon_o)$ is a q -convex exhaustion on $\mathbb{C}^n$ and, if C is sufficiently large, then

$$K \subset \{z \in \mathbb{C}^n : \psi_C(z) < 0\} \subset\subset D_i$$

If now $m = 1$, then $\tilde{q} = n$ and $D_{\varepsilon_o} = D_1 \cap D_2$, where each D_i is, in particular, $(n-1)$ -Runge in $\mathbb{C}^n$. Then the restriction map $H^{n-2}(\mathbb{C}^n, \mathcal{F}) \rightarrow H^{n-2}(D_i, \mathcal{F})$ has a dense image. This proves that $H^{n-2}(D_i, \mathcal{F}) = 0$ for $i = 1, 2$. Because $\mathbb{C}^n \setminus D_1 \cup D_2$ has no compact connected components, then, by [3], $D_1 \cup D_2$ is n -Runge in $\mathbb{C}^n$, which implies that $H^{n-1}(D_1 \cup D_2, \mathcal{F}) = 0$. Therefore from the Mayer-Vietoris sequence for cohomology

$$\begin{aligned} \dots \rightarrow H^{n-2}(D_1, \mathcal{F}) \oplus H^{n-2}(D_2, \mathcal{F}) &\rightarrow H^{n-2}(D_{\varepsilon_o}, \mathcal{F}) \rightarrow \\ &H^{n-1}(D_1 \cup D_2, \mathcal{F}) \rightarrow \dots \end{aligned}$$

it follows that $H^{n-2}(D_{\varepsilon_o}, \mathcal{F}) = 0$.

Suppose now that for any integer k with $1 \leq k \leq m$ the cohomology group $H^p(D_{i_1} \cap \dots \cap D_{i_k}, \mathcal{F}) = 0$ for all $p \geq \tilde{q} - 2$ and $i_1, \dots, i_k \in \{1, 2, \dots, m+1\}$. Then, by [5, Proposition 1], $H^{\tilde{q}-2}(D_1 \cap \dots \cap D_{m+1}, \mathcal{F}) \simeq H^{n-1}(D_1 \cup \dots \cup D_{m+1}, \mathcal{F})$. Since $\mathbb{C}^n \setminus D_1 \cup \dots \cup D_{m+1}$ has no compact connected components, then $D_1 \cup \dots \cup D_{m+1}$ is n -Runge. Therefore $H^{\tilde{q}-2}(D_1 \cap \dots \cap D_{m+1}, \mathcal{F}) = H^{n-1}(D_1 \cup \dots \cup D_{m+1}, \mathcal{F}) = 0$.

lemma 2 - *The set D_{ε_o} is not $(\tilde{q} - 1)$ -Runge in $\mathbb{C}^n$*

Proof

There exists small enough $\delta > 0$ such that the following topological sphere of real dimension $2n - m - 1$:

$$\begin{aligned} S_\delta = \{z \in \mathbb{C}^n : x_1^2 + \dots + x_m^2 + |z_{m+1}|^2 + \dots + |z_n|^2 = \delta, y_j = \\ - \sum_{i=m+1}^n + (m+1) \sum_{i=m+(j-1)(q-1)+1}^{m+j(q-1)} |z_i|^2 \text{ for } j = 1, \dots, m\} \end{aligned}$$

is contained in D_{ε_o} . It was shown by Diederich-Fornaess [4] that S_δ is not homologous to 0 in D_{ε_o} , because D_{ε_o} does not meet

$$E = \{z \in \mathbb{C}^n : x_1 = \dots = x_m = z_{m+1} = \dots = z_n = 0\}.$$

Suppose now that D_{ε_o} is $(\tilde{q} - 1)$ -Runge in $\mathbb{C}^n$. Then there exists a smooth $(\tilde{q} - 1)$ -convex exhaustion function ϕ on $\mathbb{C}^n$ such that

$$S_\delta \subset \tilde{D}_o = \{z \in \mathbb{C}^n : \phi(z) < 0\} \subset\subset D_{\varepsilon_o}.$$

Since $\tilde{D}_o \cap E = \emptyset$, then S_δ is not homologous to 0 in $\tilde{D}_o$. Moreover, there exists a sufficiently large constant $C > 0$ such that

$$B \subset \tilde{D}_C = \{z \in \mathbb{C}^n : \phi(z) < C\}$$

But since in addition ϕ is $(\tilde{q} - 1)$ -convex, it follows from [6] that

$$H_{2n-(m+1)}(\tilde{D}_o, \mathbb{R}) \simeq H_{2n-(m+1)}(\tilde{D}_C, \mathbb{R}),$$

which is false, because S_δ is clearly homologous to 0 in $\tilde{D}_C$.

References

- [1] A. Andreotti and H. Grauert. Théorèmes de finitude pour la cohomologie des espaces complexes. Bull. Soc. Math. France 90 (1962), 153-259
- [2] M. Coltoiu, On q -Runge Pairs. Ann. Scuola. Norm. Pisa Cl. Sci. (5) Vol. II (2003), pp. 231 – 235
- [3] M. Coltoiu and A. Silva, Behnke-Stein theorem on complex spaces with singularities. Nagoya Math. J. 137 (1995), 183 – 194
- [4] H. Diederich, J. E. Fornaess, Smoothing q -convex functions and vanishing theorems. Invent. Math. 82. 291-305 (1985)
- [5] K. Matsumoto, On the cohomological completeness of q -complete domains with corners. Nagoya Math. J. Vol. 165 (2002), 105 – 112
- [6] M. Peternel, Ein Lefschetz-Satz für Schnitte in projektiv-algebraischen Mannigfaltigkeiten. Math. Ann. 246, 361 – 388 (1981)